

Individual Differences in Anterior Cingulate Activation Associated with Attentional Bias Predict Cocaine Use After Treatment

Reshmi Marhe¹, Maartje Luijten^{1,2}, Ben JM van de Wetering³, Marion Smits² and Ingmar HA Franken^{*1,4}

¹Department of Psychology, Erasmus University Rotterdam, Rotterdam, The Netherlands; ²Department of Radiology, Erasmus MC—University Medical Center Rotterdam, Rotterdam, The Netherlands; ³Bouman-GGZ, Rotterdam, The Netherlands; ⁴Department of Child and Adolescent Psychiatry, Erasmus MC—University Medical Center Rotterdam, Rotterdam, The Netherlands

Drug-dependent patients often relapse into drug use after treatment. Behavioral studies show that enhanced attentional bias to drug cues is a precursor of relapse. The present functional magnetic resonance imaging (fMRI) study examined whether brain regions involved in attentional bias are predictive of cocaine use after treatment. Attentional bias-related brain activity was measured—with a cocaine Stroop task—in cocaine-dependent patients during their first week in detoxification treatment and was used to predict cocaine use at 3-month follow-up. The predictive value of attentional bias-related brain activity in *a priori* defined regions of interest, in addition to other measures such as self-reports of substance severity, craving, and behavioral attentional bias were examined. The results show that craving in the week before treatment and individual variability in attentional bias-related activity in the dorsal anterior cingulate cortex (dACC) were significant predictors of days of cocaine use at 3-month follow-up and accounted for 45% in explained variance. Brain activity in the dACC uniquely contributed 22% of explained variance to the prediction model. These findings suggest that hyperactive attentional bias-related brain activity in the dACC might be a biomarker of relapse vulnerability as early as in the first week of detoxification treatment. Ultimately, this may help to develop individually tailored treatment interventions to reduce relapse risk.

Neuropsychopharmacology (2013) **38**, 1085–1093; doi:10.1038/npp.2013.7; published online 30 January 2013

Keywords: cocaine; relapse; dorsal anterior cingulate cortex; attentional bias; functional MRI; addiction Stroop task

INTRODUCTION

The main goal of most substance dependence treatment is to prevent patients from relapsing into substance use. Despite the efforts to treat substance-dependent patients effectively, treatment dropout rates are generally > 50% and most patients subsequently relapse (Miller, 1996; Franken and Hendriks, 1999; Hättenschwiler *et al*, 2000). The identification of predictors of substance use relapse have been of high interest in addiction research (eg, Miller *et al*, 1996; Donovan, 1996; McKay, 1999). Particularly in cocaine addiction, a variety of predictors of relapse have been tested including demographic variables, substance use severity, and craving (for a review, see Poling *et al*, 2007). These predictors are largely based on self-report measures. However, over the years the idea that addiction is a relapsing brain disorder (Leshner, 1997) has gained more interest. Support for this idea comes from recent studies

suggesting that (neuro)cognitive measures might be better predictors of relapse than self-report measures (Marissen *et al*, 2006; Kosten *et al*, 2006). Currently, addiction research is focusing more on the role of cognitive predictors in relapse to substance use, and specifically the neural correlates of cognitive control (Garavan and Hester, 2007; Goldstein and Volkow, 2011).

Attentional bias is one of the most studied cognitive processes in addiction. Attentional bias in addiction refers to the automatically enhanced cognitive processing of drug-related (salient) stimuli compared with neutral (non-salient) stimuli. Recent theories suggest that it is associated with craving and therefore has an important role in the maintenance and relapse of drug dependency (Franken, 2003; Field and Cox, 2008). A widely used task to measure attentional bias is the addiction Stroop task (Cox *et al*, 2006). Behavioral studies that have used the addiction Stroop paradigm in substance-dependent individuals have indeed found an attentional bias toward the substance of abuse (for a review, see Cox *et al*, 2006). More importantly, several studies show that attentional bias to substance-related stimuli (on the behavioral level) is a predictor of treatment outcome and relapse in smoking (Waters *et al*, 2003; Powell *et al*, 2010), alcohol (Cox *et al*, 2002, 2007),

*Correspondence: Dr IHA Franken, Institute of Psychology, Erasmus University Rotterdam, P.O. Box 1738, Rotterdam 3000 DR, The Netherlands, Tel: +31 10 408 9563, Fax: +31 10 408 9009, E-mail: franken@fsw.eur.nl

Received 21 August 2012; revised 18 December 2012; accepted 20 December 2012; accepted article preview online 7 January 2013

heroin (Marissen *et al*, 2006), and cocaine use (Carpenter *et al*, 2006; Marhe *et al*, 2012). These results indicate a possible clinical relevance of attentional bias. In addition, more recent studies have taken an interest in studying the neural substrates of attentional bias in addiction.

Brain regions that have been found to be associated with attentional bias in addiction include (prefrontal) cortices such as the anterior cingulate cortex (ACC; Luijten *et al*, 2011, 2012; Nestor *et al*, 2011; Vollstädt-Klein *et al*, 2012), dorsolateral prefrontal cortex (DLPFC; Hester and Garavan, 2009; Luijten *et al*, 2012; Vollstädt-Klein *et al*, 2012), insula (Luijten *et al*, 2011; Vollstädt-Klein *et al*, 2012), and subcortical structures such as the nucleus accumbens (Nestor *et al*, 2011) and amygdala (Vollstädt-Klein *et al*, 2012; Janes *et al*, 2010a). Most pertinent to the present study is that attentional bias-related brain activity in the insula and dorsal ACC (dACC) has been found to be predictive of smoking relapse (Janes *et al*, 2010b). Involvement of the ACC in response to cocaine cues has also been found in a study with cocaine users (using the addiction Stroop task; Goldstein *et al*, 2009). However, no study to date has examined whether brain regions involved in attentional bias to cocaine stimuli (as measured by a Stroop task) are predictive of cocaine relapse after treatment.

The goal of the present study therefore was to examine whether brain activity during a cocaine Stroop task measured in the first week of detoxification treatment would be associated with cocaine use measured at 3-month follow-up. Based on aforementioned studies, we hypothesized that brain regions involved in attentional bias (ie, dACC, DLPFC, insula, nucleus accumbens, and amygdala) would be associated with cocaine use at follow-up.

MATERIALS AND METHODS

Participants

Participants were 34 cocaine-dependent inpatients recruited from an addiction treatment center (Bouman-GGZ) in Rotterdam, the Netherlands. Inclusion criteria were (1) age between 18 and 65 years; (2) DSM-IV diagnosis for cocaine dependence (assessed by both a physician and a research psychologist); and (3) the ability to speak, read, and write in Dutch at an eighth-grade literacy level. Exclusion criteria were (1) indications of severe psychopathology (ie, psychosis and severe mood disorder, as assessed by a physician); (2) self-reported color blindness or (non-corrected) defective vision; (3) pregnancy or breast-feeding; and (4) functional magnetic resonance imaging (fMRI) contra-indications. We had to exclude the data of eight participants. Reasons for exclusion were loss to follow-up ($n = 1$); mismatch between self-reported days of cocaine use and urine results at follow-up ($n = 1$); vision problems ($n = 2$); an accuracy rate on the Stroop task of $< 60\%$ ($n = 3$); these participants also showed excessive head motion of > 3 mm; major anatomical abnormality observed on MRI ($n = 1$). The final sample consisted of 26 cocaine-dependent patients (for demographics, see Table 1).

Approval of the study was received from the Ethics Committee of the Erasmus MC—University Medical Center Rotterdam, Rotterdam, the Netherlands. All procedures were carried out with the adequate understanding and

written informed consent of the participants. All participants received a financial compensation of 35 euros after completion of the fMRI measure and 25 euros after completion of the 3-month follow-up measure. The present study was part of a larger study that investigated relapse using other psychophysiological and cognitive measures (not reported in the present study).

Stroop Task

A cocaine Stroop task was used to measure attentional bias for cocaine-related stimuli. The task contained 10 cocaine words (Dutch equivalent of basepipe, crack, smoke, cocaine, blow, line, coke, snort, powder, and base), 10 neutral words drawn from the category 'indoor features' (Dutch equivalent of blanket, sofa, oven, lamp, attic, cabinet, armchair, tap, couch, and stove), and 10 letter strings (eg, MMMM; Waters *et al*, 2012). Letter strings were included to increase the number of conditions in the task which, in combination with a semi-random block order, reduced multicollinearity between parameter estimates for cocaine and neutral words (see Supplementary Information, Part I for more information on the order of task conditions). Words and letter strings were presented in the colors blue, yellow, green, or red. Participants were instructed that they should ignore the meaning of the word and only respond to the color that the word was presented in. They were instructed to indicate as rapidly and accurately as possible in which color the word was presented by pressing one of four response buttons with corresponding colors.

Table 1 Demographic and Substance Use Variables^a

Subject variable	Final sample ($n = 26$)
<i>Demographic variables</i>	
Age	38.7 (9.2)
Males (%)	85
Education (%)	
Primary education	4
Junior secondary education	54
Senior secondary education	31
Higher education	11
<i>Substance use variables</i>	
Total years of cocaine use	11.4 (6.3)
Number of days of cocaine use in 30 days before treatment entry	16.5 (11.9)
Cocaine administration route (%)	
Snorting	31
Smoking	65
Intravenous	4
<i>Craving in the week before treatment</i>	
OCDUS desire and control (1–5)	2.8 (1.1)

Abbreviation: OCDUS, Obsessive Compulsive Drug Use Scale.

^aValues are means (SD) unless otherwise indicated (%).

The task started with 32 practice trials consisting of only letter strings. Next, in the experimental phase, stimuli were presented in a blocked design with three categories (cocaine, neutral, and letter strings). Every category was presented six times, resulting in a total of 18 blocks. Each block contained 10 words. After each six blocks, there was a 38-s resting period. Order of presentation of the task conditions was semi-random; block order was the same for each participant. Each experimental trial began with a 250-ms fixation cross, followed by stimulus presentation with a duration of 1750 ms. Response could be given within this timeframe. After response, the stimulus remained on screen. No feedback regarding performance accuracy was provided to the participants.

Procedure

Patients entered the treatment center for an inpatient detoxification treatment. A standard detoxification treatment in this setting has a duration of 3 weeks. The specific goal of this treatment is to reduce cocaine withdrawal symptoms by means of psychoeducation about the detox symptoms and individual therapy based on cognitive-behavioral techniques. Afterwards, the patients start a follow-up treatment at a different department within the same treatment center. This is usually a rehabilitation program with a variable duration between 1 month and 2 years, depending on the need for treatment.

Patients that met the study criteria were informed about the study on the second day of their detoxification treatment. They had 24 h to decide whether to participate. Volunteers signed the informed consent on the third day of detoxification treatment. Cocaine use severity was assessed using the Addiction Severity Index (ASI; McLellan *et al*, 1980; Hendriks *et al*, 1989) and past week craving using the cocaine version of the Obsessive Compulsive Drug Use Scale (OCDUS; Franken *et al*, 2002). On the fourth day of treatment, participants were escorted to the Erasmus Medical Center where they performed the cocaine Stroop task in the scanner. After completion, participants were escorted back to the treatment center.

To ensure retention and compliance for the follow-up assessment, the researcher first collected multiple contact information at the start of the study. For example, we collected phone numbers and e-mail addresses of the participant but also of their family, friends, social workers, and other professionals involved with the patient. Second, the participant was informed that (s)he would receive a financial compensation of 25 euros for completion of the follow-up assessment. Third, participants were ensured that information on current use (both self-reports and urine screens) was used for research purposes only. Participants were contacted 3 months after study participation via telephone and/or e-mail to set up an appointment for a face-to-face follow-up interview. Follow-up tests were performed in the treatment center. If a participant was out of treatment at this point, then he/she was asked to return once to the treatment center to complete the follow-up tests. If the participant was unable to travel to the main treatment center for the assessment, then the researcher would perform the assessment in the (for the participant) nearest Bouman-GGZ treatment facility. If that was also not

possible, then we arranged transfer by taxi for the participant to the treatment center.

Outcome Measure

The number of days of recent cocaine use was measured 3 months after study participation. At follow-up, participants were asked to report the number of days they had used cocaine in the last 30 days, which we labeled 'recent cocaine use'. Self-reports were biochemically verified by means of urine screens. All self-reports of 0 days of use in the last 30 days were confirmed by a negative urine screen, except for one participant who reported 0 days of use while the urine screen was positive. This participant was excluded from the analysis (see Participants section). In the final sample, 11 participants reported not to have used cocaine in the last month and 15 participants reported to have used cocaine 12.1 days on average (range 1–30, SD 10.6) in the last month.

fMRI Data Acquisition and Processing

Imaging data were obtained with a 3T GE Healthcare MRI scanner. Blood oxygen level-dependent (BOLD) sensitive functional echo-planar imaging T2*-weighted images were acquired in 34 axial slices (thickness = 2.6 mm, interslice gap = 0.4 mm) covering the entire supratentorial brain with a repetition time (TR) of 2500 ms, echo time (TE) of 30 ms, field of view (FOV) of 220 mm, and matrix size of 64 × 64. A structural 3-dimensional inversion recovery (IR) fast spoiled gradient recalled echo (FSPGR) T1-weighted image was acquired in 192 axial slices (thickness = 1.6 and 0.8 mm overlap, resulting in an effective slice thickness of 0.8 mm) with TR of 12.0 ms, TE of 3.7 ms, a rectangular FOV of 250 × 175 mm², and matrix size of 416 × 256 mm. Due to an unexpected scanner shutdown, we were forced to continue the project on a different 3T scanner by the same vendor (GE Healthcare, Milwaukee, WI, USA) during the study. Data of $n = 8$ were acquired on the Signa HDxt scanner and data of $n = 18$ were acquired on the Discovery MR750 scanner. In Supplementary Information (Part VII), we describe additional analyses regarding the use of two scanners. There we report temporal signal-to-noise maps of both scanners and we conducted the main statistical analyses with scanner type included as covariate, showing that it is unlikely that the use of two scanners influenced the present findings.

Functional images were analyzed using SPM8 (Statistical Parametric Mapping; Wellcome Trust Centre for Neuroimaging, London, UK). Preprocessing of the functional data included realignment and unwarping of all functional images. Next, the anatomical scan was co-registered to the mean T2*-weighted image and subsequently segmented into gray and white matter. Segmentation parameters were used for normalization using the SPM T1 MNI 512 template. Normalized images were spatially smoothed with an 8-mm full-width at half-maximum Gaussian filter. The three task conditions, cocaine words, neutral words, and letters were modeled in the context of the general linear model, using delta functions convolved with a canonical hemodynamic response function (error rates were not associated with activation estimates, see Supplementary Information,

Part IV for results). The contrast reflecting brain activation associated with attentional bias (cocaine words minus neutral words) was calculated for each individual. Subsequently, five *a priori* regions of interest (ROIs) were selected based on their presumed role in attentional bias, including the bilateral dACC, DLPFC, insula, nucleus accumbens, and amygdala (Figure 1; see Introduction section for rationale). ROIs for the DLPFC, insula, and amygdala were derived from the automatic anatomical labeling (AAL) atlas (Tzourio-Mazoyer *et al*, 2002). Since the dACC and nucleus accumbens are not included in the AAL atlas, an 8-mm sphere around MNI coordinates ± 8 30 32 was used as an ROI for the dACC (Mars *et al*, 2005) and an 8-mm sphere around MNI coordinates ± 10 12 -2 was used for the nucleus accumbens (Knutson *et al*, 2008). To test the main attentional bias effect in the ROIs, the cocaine minus neutral words contrast was entered in a random effects one-sample *t*-test. Results were thresholded at $p < 0.05$, Family Wise Error (FWE) corrected for multiple comparisons across the search volume (small volume correction: Friston *et al*, 1996; Worsley *et al*, 1996). To do so, analyses were first thresholded at $p < 0.001$ uncorrected with 20 contingently activated voxels (160 mm^3 ; see Supplementary Information, Part V), and then corrected using a small volume correction ($p < 0.05$ FWE corrected) in which the search volume was restricted by the *a priori* defined ROIs.

Statistical Analyses

First, to demonstrate attentional bias on a behavioral level for cocaine words, behavioral data of the Stroop task were examined (means are displayed in Table 2). Differences in reaction times (RTs) and accuracy scores for the cocaine words and the neutral words were tested using a paired samples *t*-test. To use the Stroop effect as a predictor in regression models (described below), a single differential

score was calculated for RT and for accuracy (by subtracting results on cocaine words from neutral words).

Second, linear regression analysis was used to examine the predictive value of multiple variables (ie, addiction severity, craving, behavioral and brain-related attentional bias) in the prediction of the outcome measure 'recent cocaine use'. Due to the relative large amount of predictor variables, a stepwise regression analysis was performed (Field, 2009). The stepwise method tests, at each addition of a variable into the regression model, which variable contributes the least to the prediction of recent cocaine use and removes this variable from the model. By using this method all redundant predictors are filtered out of the model. Because we were interested in the unique contribution of three clusters of variables (ie, self-reports, behavior, and imaging) we performed the regression analysis in three steps. Step 1 contained four self-report variables: total years of cocaine use, number of days of cocaine use in 30 days before treatment entry, and cocaine administration route (ie, addiction severity) as well as craving in the week before treatment. Since cocaine administration route is a categorical variable with three levels, two dummy variables were created. Step 2 contained two variables of behavioral data: Stroop effect RT and Stroop effect accuracy. Step 3 contained the imaging variables (extracted contrast values for cocaine minus neutral words in the five bilateral ROIs; left and right dACC, left and right DLPFC, left and right insula, left and right nucleus accumbens, left and right amygdala). In each step, the selection criteria for inclusion and exclusion of predictors in the model were $F_{\text{enter}}: p \leq 0.05$ and $F_{\text{remove}}: p \geq 0.10$.

Third, we wanted to check the robustness of the individual predictors that were included in the model based on the results of the stepwise regression. To do so, bootstrapping was used with 2000 bootstrapped samples and a 95% confidence interval (CI; Efron and Tibshirani, 1993; Mooney and Duval, 1993). The bootstrap procedure




Figure 1 *A priori* defined, literature-based regions of interest involved in attentional bias to substance cues. dACC, dorsal anterior cingulate cortex; DLPFC, dorsolateral prefrontal cortex; NACC, nucleus accumbens.

can only be conducted when predictor variables are forcedly entered into the regression model (instead of stepwise). Therefore, predictors that met the inclusion criteria of the stepwise method were entered in a hierarchical regression analysis.

Finally, predictors were checked for multicollinearity by means of tolerance statistics. All statistical analysis were conducted with SPSS 18.0 (IBM Corporation, Armonk, NY, USA).

RESULTS

Attentional Bias

We found a significant difference between Stroop RTs on cocaine words and neutral words, ($t[25] = 6.87, p < 0.001$), with cocaine-dependent patients responding slower to cocaine-related stimuli than to neutral stimuli. This indicates the presence of an attentional bias. No differences in accuracy between cocaine and neutral words were found, ($t[25] = -1.84, ns$).

For the imaging data, we did not find significant activation ($p < 0.05$ FWE small volume corrected) for the cocaine minus neutral words contrast in any of the ROIs within the patient sample.

Predictors of Recent Cocaine Use

The stepwise regression yielded that two predictors—craving in the week before treatment and attentional bias-related activity in the right dACC—met the selection criterion for inclusion in the model. Other self-report, behavioral, and imaging variables were excluded based on the criterion (see Table 3a). Model statistics are displayed in Table 3b. The first step, including craving in the week before treatment, accounted for 23% of explained variance in recent cocaine use. Individual differences in brain activation for the attentional bias contrast in the right dACC added another 22% explained variance to the model. Thus, together both predictors accounted for 45% of explained variance in the prediction of recent cocaine use.

Bootstrapping confirmed that both individual predictors were robust predictors of treatment outcome (see Table 3c). The coefficients indicated a positive association between the predictors and the outcome measure. These findings showed that higher self-reported past-week craving and enhanced activation for cocaine words vs neutral words in the right dACC (both measured during the first week of treatment) were associated with more days of recent cocaine

use (measured at 3-month follow-up). Direct associations between each significant predictor and recent cocaine use are displayed in Figure 2.

Finally, collinearity statistics indicated that the predictors were not associated with each other (tolerance = 0.98).

Table 3a Statistics of Predictor Variables Added in the Stepwise Regression Analysis

Predictor variables	Standardized coefficients (= β)	t statistic	p-Value
<i>Self-reports</i>			
Total years of cocaine use	0.14	0.88	0.387
Number of days of cocaine use in 30 days before treatment entry	0.04	0.22	0.831
Cocaine administration route			
Snorting vs smoking	0.18	1.09	0.287
Snorting vs intravenous	0.08	0.46	0.647
Craving in the week before treatment	0.41	2.57	0.017 ^a
<i>Behavior</i>			
Stroop effect RT	-0.24	-1.51	0.145
Stroop effect accuracy	-0.11	-0.66	0.519
<i>Imaging^b</i>			
l-dACC	-0.53	-1.70	0.104
r-dACC	0.47	3.00	0.006 ^a
l-DLPFC	-0.35	-1.39	0.178
r-DLPFC	-0.27	-1.01	0.322
l-insula	-0.36	-1.41	0.174
r-insula	-0.06	0.21	0.832
l-nucleus accumbens	-0.11	-0.40	0.694
r-nucleus accumbens	-0.05	-0.17	0.868
l-amygdala	-0.37	-1.53	0.140
r-amygdala	0.04	0.15	0.884

Abbreviations: RT, reaction time; l, left; r, right; dACC, dorsal anterior cingulate cortex; DLPFC, dorsolateral prefrontal cortex.

^aMeet selection criteria $F_{enter} : p \leq 0.05$ and $F_{remove} : p \geq 0.10$.

^bAttentional bias contrast (cocaine minus neutral).

Table 3b Model Statistics and Change Statistics of the Final Model Predicting Recent Cocaine Use

Predictor variables	Model statistics			Change statistics		
	R ²	F Statistic	p-Value	ΔR^2	ΔF statistic	Δp -Value
Step 1	0.23	7.15	0.013			
Craving in the week before treatment						
Step 2	0.45	9.26	0.001	0.22	9.00	0.006
r-dACC ^a						

Abbreviations: dACC, dorsal anterior cingulate cortex; r, right.

^aAttentional bias contrast (cocaine minus neutral).

Table 2 Mean Reaction Times (milliseconds) and Accuracy Scores (percentages) for the Stroop Task

	Mean	SD
RT cocaine words	972	137
RT neutral words	886	126
Accuracy cocaine words	91	8
Accuracy neutral words	93	8

Abbreviation: RT, reaction time.

DISCUSSION

The present study is the first to examine to what extent attentional bias-related brain activity predicts recent cocaine use after treatment, over and above other relevant predictors such as craving, substance use severity, and behavioral measures of attentional bias. We found that both higher levels of self-reported craving in the week before treatment and enhanced attentional bias-related brain activity in the right dACC—measured in the first week of detoxification treatment—were associated with cocaine use at 3-month follow-up. Cocaine use severity, behavioral measures of attentional bias, and other brain regions that have previously been shown to be associated with attentional bias (ie, DLPFC, insula, nucleus accumbens, and amygdala) were not significantly associated with cocaine use at 3-month follow-up. In addition, we found that when controlling for self-reported craving, individual differences in right dACC activity made a rather large and unique contribution to the prediction of recent cocaine use. These findings suggest that a neural correlate of attentional bias in the dACC might be a useful biomarker of cocaine use after treatment. Most importantly, as early as in the first week of treatment our findings provide information on an individual's risk to use cocaine again 3 months after treatment.

The dorsal part of the ACC has been implicated in cognitive processes such as salience detection (Seeley *et al*, 2007), conflict monitoring in general (Botvinick *et al*, 2004; Kerns *et al*, 2004; Egner *et al*, 2008) and in the presence of emotionally salient distractors (specifically the right dACC; Haas *et al*, 2006). Hyperactivity in the ACC associated with attentional bias to substance-related cues has previously been shown in smokers (Luijten *et al*, 2011, 2012; Nestor

et al, 2011; Janes *et al*, 2010b) and alcohol-dependent patients (Vollstädt-Klein *et al*, 2012), and is thought to reflect enhanced conflict when task performance is being interfered by the automatic detection of salient substance-related stimuli. However, the present design does not rule out alternative interpretations of the dACC involvement. For example, it might be possible that dACC activity is related to the greater requirement for cognitive control during the drug trials, rather than the content of the stimuli *per se*.

In relation to relapse, it is suggested that individual variations in the brain's ability to control cognitive conflict may account for differences in risk of relapse (Garavan and Hester, 2007). Our results support this idea by finding that when more top-down control is necessary (ie, increased dACC activity) to focus on a cognitive task in the presence of distracting drug cues one is more vulnerable to use cocaine after treatment. Additionally, individuals who need less top-down control (as reflected by less to no attentional bias-related dACC activity) are less susceptible to return to daily use of cocaine. This is in line with findings of Janes *et al* (2010b) who found that attentional bias-related dACC activity was predictive of relapse in smokers.

Concerning the self-report measures we examined in the current study, we found that self-reported craving in the week before treatment—but not substance use severity—was predictive of cocaine use at 3-month follow-up. Craving constitutes a central role in established theories of addiction (Robinson and Berridge, 1993). Indeed several studies have reported that subjective craving is associated with relapse and treatment outcome in cocaine-dependent patients (Weiss *et al*, 2003; Paliwal *et al*, 2008; Preston *et al*, 2009). More recent models of addiction propose a role for both craving and attentional bias in the maintenance of substance use and relapse (Franken, 2003; Field and Cox, 2008). Our results support this by finding that both self-reported craving in the week before treatment and individual variability in dACC activity related to attentional bias were associated with cocaine use at follow-up, suggesting a role for both processes in relapse. However, these models suggest that attentional bias predicts relapse via craving and not directly, whereas our results showed that dACC activity was a strong predictor, even when controlling for self-reported craving. This suggests that enhanced attentional bias-related brain activity might be independently associated with drug relapse.

Somewhat unexpected in the light of the recent discussion on the relation between craving and attentional bias (Field,

Table 3c Bootstrap Results for Individual Predictors in the Final Model Predicting Recent Cocaine Use

Predictor variables	Unstandardized coefficients ($=\beta$)	Standard error (=SE)	Bootstrapped 95% confidence interval
Craving in the week before treatment	3.82	1.48	1.48–7.36 ^a
r-dACC ^b	15.62	5.35	4.61–25.14 ^a

Abbreviations: r, right; dACC, dorsal anterior cingulate cortex.

^aSignificant (value 0 is not in the confidence interval).

^bAttentional bias contrast (cocaine minus neutral).




Figure 2 Correlation between (a) craving in the week before treatment and recent cocaine use and (b) attentional bias-related activation in the right dACC and recent cocaine use. dACC, dorsal anterior cingulate cortex.

Munafò and Franken, 2009), no significant correlations between craving, attentional bias, and dACC activation were observed (see Supplementary Information, Part II for results). The meta-analysis by Field *et al* (2009) showed that subjective craving is consistently (though weakly) correlated with behavioral and EEG indices of attentional bias. Concerning the behavioral measures, it is possible that the statistical power of the present study was too low to yield significant results given the fact that the correlations between craving and the Stroop task are known to be weak ($r = 0.15$). In addition, dACC activation was not found to be associated with craving nor behavioral attentional bias. This is somewhat unexpected since all measures could be regarded as measures of the same underlying processing: 'motivational salience'. Although exact interpretation is not straightforward (also given the relative small n), it seems at least to be an indication that the ACC does not necessarily be the neurobiological structure responsible for cocaine craving. Further, it should be kept in mind that craving was not measured during fMRI scanning but was an average measure (OCDUS) of the craving in the week before the fMRI measurement.

The present study indicates that cocaine use after treatment is better predicted by neuroimaging than behavioral measures of the addiction Stroop effect. Other fMRI studies examining neurocognitive predictors of substance relapse have reported similar results in methamphetamine relapse (Paulus *et al*, 2005) and cocaine relapse (Brewer *et al*, 2008; Jia *et al*, 2011). Although previous studies have reported an association between attentional bias on the behavioral level and drug relapse (Marissen *et al*, 2006; Carpenter *et al*, 2006) we could not find any significant associations between behavioral measures and cocaine use outcome (see Supplementary Information, Part III for results). It might be that functional neuroimaging is a better predictor because it is a more sensitive measure of individual differences in biased attention to drug cues than behavioral measures.

Possible clinical implications of our results include the use of dopaminergic manipulation to reduce attentional bias-related brain activity (Ersche *et al*, 2010; Goldstein *et al*, 2010; Luijten *et al*, 2012) or attentional retraining (Schoenmakers *et al*, 2010). Patients who are vulnerable to relapse due to enhanced attentional bias for drug cues might benefit from such approaches, although these interventions need further investigation. Future intervention studies using either psychopharmacological interventions or cognitive training may obtain valuable information by identifying those patients with higher levels of dACC activity in response to drug cues.

The present study had limitations. First, we only measured days of cocaine use in the last month of the 3-month follow-up period. We could not rule out that patients who reported to have used 0 days in the last month did not use in the 2 months before follow-up and could therefore not term them as 'non-relapsers'. Thus, with the present design it was not possible to predict relapse (yes *vs* no) which makes it difficult to compare the present results with other fMRI studies on relapse prediction (eg, Janes *et al*, 2010b; Paulus *et al*, 2005). However, it has been found that continuous and dichotomous drug use outcome measures are strongly related to each other (McKay *et al*, 2001),

suggesting that the present outcome measure does provide some information on relapse risk. To examine associations between study variables and different drug use outcome measures, future studies should use multiple outcome measures (eg, time to relapse, relapse *vs* non-relapse; see for a discussion also Miller *et al*, 1996). Second, while we found an association between enhanced attentional bias-related brain activity and cocaine use outcome in patients after treatment, we could not demonstrate an overall within-group attentional bias effect in the ROIs. The absence of an overall effect on the attentional bias contrast (cocaine minus neutral words) is not an unusual finding in cocaine Stroop fMRI research (Goldstein *et al*, 2007, 2009, 2010; Moeller *et al*, 2012) and may be due to the presence of individual differences in brain activity related to attentional bias (Hester and Garavan, 2009), as manifested by the fact that brain activity was predictive of relapse in a population of variable relapse risk. We would like to emphasize, however, that the task did activate regions that are typically expected to be activated for a Stroop task such as the dACC and the DLPFC for the neutral and cocaine condition separately (see Supplementary Figure 1 in Supplementary Information, Part VI). Third, we did not assess other psychiatric disorders using a structured interview or questionnaire and therefore could not examine the influence of these disorders on cocaine use outcome. Finally, we had a large number of predictors and a relatively small sample which could have led to over-fitting of the prediction model. Therefore, we performed a stepwise regression analysis to remove all redundant predictors. To reduce the probability of type II errors, we included variables in a hierarchical manner based upon previous literature (self-report measures first, behavior measures second, and neuroimaging measures third) so that the predictors are evaluated in each step separately.

In conclusion, the current study showed the clinical relevance of measuring attentional bias-related activity in the dACC in cocaine-dependent patients during the starting phase of treatment. Not only does this provide valuable information about the neurocognitive mechanisms of addiction and of addiction treatment processes, these measures may also help us identifying individuals who are at risk of relapse into cocaine use after detoxification treatment. Although direct clinical implications are not very easy to implement using fMRI methodology, ultimately findings like these may help to develop clinical profiles based on neurocognitive measures to design tailor-made follow-up treatment schedules. Future treatment intervention studies should investigate how this information might benefit relapse-vulnerable patients.

ACKNOWLEDGEMENTS

We thank Radha Jagroep, Yavuz Kilic, and Alexandra de Raaij for their assistance with data collection. This study was supported by a grant of the Netherlands Organization for Health Research and Development (ZonMw Grant no. 31160203 to IHA Franken). The funding organization had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

DISCLOSURE

The authors declare no conflict of interest.

REFERENCES

- Botvinick MM, Cohen JD, Carter CS (2004). Conflict monitoring and anterior cingulate cortex: an update. *Trends Cogn Sci (Regul Ed)* 8: 539–546.
- Brewer JA, Worhunsky PD, Carroll KM, Rounsaville BJ, Potenza MN (2008). Pretreatment brain activation during stroop task is associated with outcomes in cocaine-dependent patients. *Biol Psychiatry* 64: 998–1004.
- Carpenter KM, Schreiber E, Church S, McDowell D (2006). Drug Stroop performance: relationships with primary substance of use and treatment outcome in a drug-dependent outpatient sample. *Addict Behav* 31: 174–181.
- Cox WM, Fardari JS, Pothos EM (2006). The addiction-stroop test: theoretical considerations and procedural recommendations. *Psychol Bull* 132: 443–476.
- Cox WM, Hogan LM, Kristian MR, Race JH (2002). Alcohol attentional bias as a predictor of alcohol abusers' treatment outcome. *Drug Alcohol Depend* 68: 237–243.
- Cox WM, Pothos EM, Hosier SG (2007). Cognitive-motivational predictors of excessive drinkers' success in changing. *Psychopharmacology (Berl)* 192: 499–510.
- Donovan DM (1996). Assessment issues and domains in the prediction of relapse. *Addiction* 91: S29–S36.
- Efron B, Tibshirani RJ (1993). *An Introduction to the Bootstrap*. Chapman & Hall: New York, NY.
- Egner T, Etkin A, Gale S, Hirsch J (2008). Dissociable neural systems resolve conflict from emotional versus nonemotional distracters. *Cereb Cortex* 18: 1475–1484.
- Ersche KD, Bullmore ET, Craig KJ, Shabbir SS, Abbott S, Müller U et al (2010). Influence of compulsivity of drug abuse on dopaminergic modulation of attentional bias in stimulant dependence. *Arch Gen Psychiatry* 67: 632–644.
- Field A (2009). *Discovering Statistics Using SPSS*. 3rd edn. Sage: London, UK.
- Field M, Cox WM (2008). Attentional bias in addictive behaviors: a review of its development, causes, and consequences. *Drug Alcohol Depend* 97: 1–20.
- Field M, Munafo MR, Franken IHA (2009). A meta-analytic investigation of the relationship between attentional bias and subjective craving in substance abuse. *Psychol Bull* 135: 589–607.
- Franken IHA (2003). Drug craving and addiction: integrating psychological and neuropsychopharmacological approaches. *Prog Neuropsychopharmacol Biol Psychiatry* 27: 563–579.
- Franken IHA, Hendriks VM (1999). Predicting outcome of inpatient detoxification of substance abusers. *Psychiatr Serv* 50: 813–817.
- Franken IHA, Hendriks VM, van den Brink W (2002). Initial validation of two opiate craving questionnaires—the obsessive compulsive drug use scale and the desires for drug questionnaire. *Addict Behav* 27: 675–685.
- Friston KJ, Holmes A, Poline JB, Price CJ, Frith CD (1996). Detecting activations in PET and fMRI: levels of inference and power. *Neuroimage* 4: 223–235.
- Garavan H, Hester R (2007). The role of cognitive control in cocaine dependence. *Neuropsychol Rev* 17: 337–345.
- Goldstein RZ, Alia-Klein N, Tomasi D, Carrillo JH, Maloney T, Woicik PA et al (2009). Anterior cingulate cortex hypoactivations to an emotionally salient task in cocaine addiction. *Proc Natl Acad Sci USA* 106: 9453–9458.
- Goldstein RZ, Tomasi D, Rajaram S, Cottone LA, Zhang L, Maloney T et al (2007). Role of the anterior cingulate and medial orbitofrontal cortex in processing drug cues in cocaine addiction. *Neuroscience* 144: 1153–1159.
- Goldstein RZ, Volkow ND (2011). Dysfunction of the prefrontal cortex in addiction: neuroimaging findings and clinical implications. *Nat Rev Neurosci* 12: 652–669.
- Goldstein RZ, Woicik PA, Maloney T, Tomasi D, Alia-Klein N, Shan J et al (2010). Oral methylphenidate normalizes cingulate activity in cocaine addiction during a salient cognitive task. *Proc Natl Acad Sci USA* 107: 16667–16672.
- Haas BW, Omura K, Constable RT, Canli T (2006). Interference produced by emotional conflict associated with anterior cingulate activation. *Cogn Affect Behav Neurosci* 6: 152–156.
- Hättenschwiler J, Ruesch P, Hell D (2000). Effectiveness of inpatient drug detoxification: links between process and outcome variables. *Eur Addict Res* 6: 123–131.
- Hendriks VM, Kaplan CD, Van Limbeek J, Geerlings P (1989). The addiction severity index—reliability and validity in a dutch addict population. *J Subst Abuse Treat* 6: 133–141.
- Hester R, Garavan H (2009). Neural mechanisms underlying drug-related cue distraction in active cocaine users. *Pharmacol Biochem Behav* 93: 270–277.
- Janes AC, Pizzagalli DA, Richardt S, Frederick Bde B, Holmes AJ, Sousa J et al (2010a). Neural substrates of attentional bias for smoking-related cues: an fMRI study. *Neuropsychopharmacology* 35: 2339–2345.
- Janes AC, Pizzagalli DA, Richardt S, Bd Frederick, Chuzi S, Pachas G et al (2010b). Brain reactivity to smoking cues prior to smoking cessation predicts ability to maintain tobacco abstinence. *Biol Psychiatry* 67: 722–729.
- Jia Z, Worhunsky PD, Carroll KM, Rounsaville BJ, Stevens MC, Pearlson GD et al (2011). An initial study of neural responses to monetary incentives as related to treatment outcome in cocaine dependence. *Biol Psychiatry* 70: 553–560.
- Kerns JG, Cohen JD, MacDonald AW, Cho RY, Stenger VA, Carter CS (2004). Anterior cingulate conflict monitoring and adjustments in control. *Science* 303: 1023–1026.
- Knutson B, Wimmer GE, Rick S, Hollon NG, Prelec D, Loewenstein G (2008). Neural antecedents of the endowment effect. *Neuron* 58: 814–822.
- Kosten TR, Scanley BE, Tucker KA, Oliveto A, Prince C, Sinha R et al (2006). Cue-induced brain activity changes and relapse in cocaine-dependent patients. *Neuropsychopharmacology* 31: 644–650.
- Leshner AI (1997). Addiction is a brain disease, and it matters. *Science* 278: 45–47.
- Luijten M, Veltman DJ, Hester R, Smits M, Peplinkhuizen L, Franken IHA (2012). Brain activation associated with attentional bias in smokers is modulated by a dopamine antagonist. *Neuropsychopharmacology* 37: 2772–2779.
- Luijten M, Veltman DJ, van den Brink W, Hester R, Field M, Smits M et al (2011). Neurobiological substrate of smoking-related attentional bias. *Neuroimage* 54: 2374–2381.
- Marhe R, Waters AJ, Van de Wetering BJM, Franken IHA (2012). Implicit and explicit drug-related cognitions during detoxification treatment are associated with drug relapse: an ecological momentary assessment study. *J Consult Clin Psychol* (in press).
- Marissen MAE, Franken IHA, Waters AJ, Blanken P, van den Brink W, Hendriks VM (2006). Attentional bias predicts heroin relapse following treatment. *Addiction* 101: 1306–1312.
- Mars RB, Coles MGH, Grol MJ, Holroyd CB, Nieuwenhuis S, Hulstijn W et al (2005). Neural dynamics of error processing in medial frontal cortex. *Neuroimage* 28: 1007–1013.
- McKay JR (1999). Studies of factors in relapse to alcohol, drug and nicotine use: a critical review of methodologies and findings. *J Stud Alcohol* 60: 566–576.
- McKay JR, Alterman AI, Koppenhaver J, Mulvaney F, Bovasso G, Ward K (2001). Continuous, categorical, and time to event cocaine use outcome variables: degree of intercorrelation and sensitivity to treatment group differences. *Drug Alcohol Depend* 62: 19–30.

- McLellan AT, Luborsky L, Woody GE, O'Brien CP (1980). An improved diagnostic evaluation instrument for substance abuse patients. The Addiction Severity Index. *J Nerv Ment Dis* **168**: 26–33.
- Miller WR (1996). What is a relapse? Fifty ways to leave the wagon. *Addiction* **91**: S15–S27.
- Miller WR, Westerberg VS, Harris RJ, Tonigan JS (1996). What predicts relapse? Prospective testing of antecedent models. *Addiction* **91**: S155–S172.
- Moeller SJ, Tomasi D, Woicik PA, Maloney T, Alia-Klein N, Honorio J et al (2012). Enhanced midbrain response at 6-month follow-up in cocaine addiction, association with reduced drug-related choice. *Addict Biol* **17**: 1013–1025.
- Mooney CZ, Duval RD (1993). *Bootstrapping: A Nonparametric Approach to Statistical Inference*. Sage: Newbury Park, CA.
- Nestor L, McCabe E, Jones J, Clancy L, Garavan H (2011). Differences in 'bottom-up' and 'top-down' neural activity in current and former cigarette smokers: evidence for neural substrates which may promote nicotine abstinence through increased cognitive control. *Neuroimage* **56**: 2258–2275.
- Paliwal P, Hyman SM, Sinha R (2008). Craving predicts time to cocaine relapse: further validation of the Now and Brief versions of the cocaine craving questionnaire. *Drug Alcohol Depend* **93**: 252–259.
- Paulus MP, Tapert SF, Schuckit MA (2005). Neural activation patterns of methamphetamine-dependent subjects during decision making predict relapse. *Arch Gen Psychiatry* **62**: 761–768.
- Poling J, Kosten TR, Sofuoglu M (2007). Treatment outcome predictors for cocaine dependence. *Am J Drug Alcohol Abuse* **33**: 191–206.
- Powell J, Dawkins L, West R, Powell J, Pickering A (2010). Relapse to smoking during unaided cessation: clinical, cognitive and motivational predictors. *Psychopharmacology (Berl)* **212**: 537–549.
- Preston KL, Vahabzadeh M, Schmittner J, Lin J-, Gorelick DA, Epstein DH (2009). Cocaine craving and use during daily life. *Psychopharmacology (Berl)* **207**: 291–301.
- Robinson TE, Berridge KC (1993). The neural basis of drug craving: an incentive-sensitization theory of addiction. *Brain Res Rev* **18**: 247–291.
- Schoenmakers TM, de Bruin M, Lux IF, Goertz AG, Van Kerkhof DH, Wiers RW (2010). Clinical effectiveness of attentional bias modification training in abstinent alcoholic patients. *Drug Alcohol Depend* **109**: 30–36.
- Seeley WW, Menon V, Schatzberg AF, Keller J, Glover GH, Kenna H et al (2007). Dissociable intrinsic connectivity networks for salience processing and executive control. *J Neurosci* **27**: 2349–2356.
- Tzourio-Mazoyer N, Landeau B, Papathanassiou D, Crivello F, Etard O, Delcroix N et al (2002). Automated anatomical labeling of activations in SPM using a macroscopic anatomical parcellation of the MNI MRI single-subject brain. *Neuroimage* **15**: 273–289.
- Vollstädt-Klein S, Loeber S, Richter A, Kirsch M, Bach P, von der Goltz C et al (2012). Validating incentive salience with functional magnetic resonance imaging: association between mesolimbic cue reactivity and attentional bias in alcohol-dependent patients. *Addict Biol* **17**: 807–816.
- Waters AJ, Marhe R, Franken IHA (2012). Attentional bias to drug cues is elevated before and during temptations to use heroin and cocaine. *Psychopharmacology (Berl)* **219**: 909–921.
- Waters AJ, Shiffman S, Sayette MA, Paty JA, Gwaltney CJ, Balabanis MH (2003). Attentional bias predicts outcome in smoking cessation. *Health Psychol* **22**: 378–387.
- Weiss RD, Griffin ML, Mazurick C, Berkman B, Gastfriend DR, Frank A et al (2003). The relationship between cocaine craving, psychosocial treatment, and subsequent cocaine use. *Am J Psychiatry* **160**: 1320–1325.
- Worsley KJ, Marrett S, Neelin P, Vandal AC, Friston KJ, Evans AC (1996). A unified statistical approach for determining significant signals in images of cerebral activation. *Hum Brain Mapp* **4**: 58–73.

Supplementary Information accompanies the paper on the Neuropsychopharmacology website (<http://www.nature.com/npp>)